VOL. 589, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY JANUARY 10, 1992

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CRNS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

Publisher's note V

REVIEW

Reaction kinetics processes in modern liquid chromatographic reactors
by C.-Y. Jeng and S. H. Langer (Madison, WI, USA) (Received August 20th, 1991) 1

REGULAR PAPERS

Column Liquid Chromatography

Feasibility study for the construction of an integrated expert system in high-performance liquid chromatography
by T. Hamoir, M. De Smet, H. Pyrins, P. Conti, N. Vanden Driessche and D. L. Massart (Brussels, Belgium), F. Maris and H. Hindriks (Oss, Netherlands) and P. J. Schoenmakers (Eindhoven, Netherlands) (Received June 4th, 1991) . . 31

Effect of superfluous remote polar functionality on chiral recognition
by W. H. Pirkle and C. J. Welch (Urbana, IL, USA) (Received September 2nd, 1991) 45

Silica-bonded chiral stationary phases with structurally simple π-donor chiral selectors for high-performance liquid chromatography
by L. Oliveros and C. Minguillón (Paris, France) and B. Desmazières and P.-L. Desbène (Evreux, France) (Received September 3rd, 1991) 53

Simple test for determination of the degree of distortion of the liquid-phase flow profile in columns for preparative liquid chromatography
by M. Kamiński (Gdańsk, Poland) (Received July 16th, 1991) 61

Predictability of chromatographic protein separations. Study of size-exclusion media with narrow particle size distributions
by A. M. Athalye, S. J. Gibbs and E. N. Lightfoot (Madison, WI, USA) (Received August 23rd, 1991) 71

Phosphorylated polystyrene resins in high-performance ion-exchange chromatography
by D. Letourneur, V. Migonney, D. Muller and M. Jozefowicz (Villetaneuse, France) (Received August 6th, 1991) . . 87

Cellulose beads: a weak affinity support
by M. Baeseler and H.-F. Boeden (Berlin, Germany) and R. Koelsch and J. Lasch (Halle/Saale, Germany) (Received August 14th, 1991) 93

Activation of matrices by 4,6-diphenylthieno[3,4-*d*]-1,3-dioxol-2-one-5,5-dioxide. High-performance liquid affinity chromatographic separations
by M. Hill and B. Arrio (Orsay, France) (Received August 26th, 1991) 101

Determination of chloropropanols in protein hydrolysates
by C. A. Van Bergen, P. D. Collier, D. D. O. Cromie, R. A. Lucas, H. D. Preston and D. J. Sissons (Sharnbrook, UK) (Received July 18th, 1991) 109

Purification and determination of glutamine synthetase by high-performance immunoaffinity chromatography
by J. Alhama, J. López-Barea, F. Toribio and J. M. Roldán (Córdoba, Spain) (Received September 6th, 1991) . . . 121

Affinity partitioning of erythrocytic phosphofructokinase in aqueous two-phase systems containing poly(ethylene glycol)-bound Cibacron Blue
by M. C. Tejedor, C. Delgado, M. Grupeli and J. Luque (Madrid, Spain) (Received July 1st, 1991) 127

Determination of profiles of non-collagenous proteins from rat bones by sodium dodecyl sulfate high-performance liquid chromatography
by M. E. Kunkel, A. Seo and R. L. Thomas (Clemson, SC, USA) (Received June 28th, 1991) 135

(Continued overleaf)

Contents (continued)

Determination of aflatoxin M1 using a dialysis-based immunoaffinity sample pretreatment system coupled on-line to liquid chromatography. Reusable immunoaffinity columns
by A. Farjam, N. C. van de Merbel, A. A. Nieman, H. Lingeman and U. A. Th. Brinkman (Amsterdam, Netherlands) (Received August 9th, 1991) 141

Optimization of an ion-exchange high-performance liquid chromatographic method for the determination of carboxylic acids, sugars, glycerol and ethanol in wines
by M. Calull, E. López, R. M. Marcé, J. C. Olucha and F. Borrull (Tarragona, Spain) (Received June 18th, 1991) . . 151

High-performance liquid chromatographic determination of pilocarpine hydrochloride and its degradation products using a β-cyclodextrin column
by K. D. Sternitzke, T. Y. Fan and D. L. Dunn (Fort Worth, TX, USA) (Received August 21st, 1991) 159

Separation of soy lecithin using gel permeation chromatography
by R. E. Shansky and R. E. Kane (Ridgefield, CT, USA) (Received June 25th, 1991) 165

Optimization of an isocratic high-performance liquid chromatographic separation of carotenoids
N. E. Craft and S. A. Wise (Gaithersburg, MD, USA) and J. H. Soares, Jr. (College Park, MD, USA) (Received September 10th, 1991) 171

Determination of aromatic sulfonic acids in aqueous environmental samples by anion-exchange chromatography coupled to particle beam mass spectrometry and UV spectrophotometry
by I. S. Kim, F. I. Sasinos, D. K. Rishi, R. D. Stephens and M. A. Brown (Berkeley, CA, USA) (Received September 3rd, 1991) 177

Measurement of stability constants for complexes of divalent cations with tartaric acid by ion chromatography
by F. H.-J. Lin (Tainen, Taiwan) and Cs. Horváth (New Haven, CT, USA) (Received July 30th, 1991) 185

Carbonate eluents in ion chromatography. Fractions of resin capacity linked to anions present in the eluent
by Ü. Haldna (Tallinn, Estonia) (Received July 19th, 1991) 197

Mixed potential response at metallic electrodes in potentiometric detection for ion chromatography
by P. W. Alexander (Kensington, NSW, Australia), B. K. Głód (Warsaw, Poland) and P. R. Haddad (Kensington, NSW, Australia) (Received August 15th, 1991) 201

Potentiometric detection of carboxylic acids and inorganic anions in ion-exclusion chromatography using camphorsulphonic acid as eluent
by B. K. Głód, P. R. Haddad and P. W. Alexander (Kensington, NSW, Australia) (Received September 7th, 1991) . . 209

Gas Chromatography

Accuracy of computer-simulated gas chromatographic separations based on a linear elution strength model
by M. C. Roman (Kingston, RI, USA) and R. W. Siergiej (Spring House, PA, USA) (Received August 13th, 1991) . 215

Prediction of the elution profiles of high-concentration bands in gas chromatography
by J. Roles and G. Guiochon (Knoxville and Oak Ridge, TN, USA) (Received July 10th, 1991) 223

Prediction of retention indexes. IV. Chain branching in alkylbenzene isomers with C_{10-13} alkyl chains identified in a scintillator solvent
by C. T. Peng, R. L. Hua and D. Maltby (San Francisco, CA, USA) (Received September 2nd, 1991) 231

Flame infrared emission detection of hydrogen
by M. Mofidi, M. K. Hudson and K. Underhill (Little Rock, AR, USA) (Received September 10th, 1991) 241

Gas chromatographic separation of diastereomeric amino acid derivatives on chiral stationary phases. Application to the determination of enantiomeric composition in (*S*)-(+)-2-butanol
by D.-M. Liu (Nanjing, China) (Received July 26th, 1991) 249

Application and chiral recognition of heptakis (2,6-di-O-methyl-3-O-trifluoroacetyl)-β-cyclodextrin as a stationary phase for the gas chromatographic separation of enantiomers
by G. Stoev (Sofia, Bulgaria) (Received August 12th, 1991) 257

Comparison of extraction methods for polychlorinated dibenzo-*p*-dioxins and dibenzofurans in fly ash using gas chromatography–mass spectrometry
by A. Beard, K. Naikwadi and F. W. Karasek (Waterloo, Canada) (Received August 23rd, 1991) 265

Determination of sulfur components in light petroleum streams by high-resolution gas chromatography with chemiluminescence detection
by B. Chawla and F. Di Sanzo (Paulsboro, NJ, USA) (Received July 25th, 1991) 271

Use of oxidative degradation followed by capillary gas chromatography–mass spectrometry and multi-dimensional scaling analysis to fingerprint unresolved complex mixtures of hydrocarbons
by A. T. Revill, M. R. Carr and S. J. Rowland (Plymouth, UK) (Received August 27th, 1991) 281

Determination of chemical warfare agents, their hydrolysis products and related compounds in soil
by P. A. D'Agostino and L. R. Provost (Medicine Hat, Canada) (Received August 19th, 1991) 287

Supercritical Fluid Chromatography

Analysis of benzyloxy-terminated poly(1,3,6-trioxocane)s by supercritical fluid chromatography
by H. Pasch, H. Krüger, H. Much and U. Just (Berlin, Germany) 295

Planar Chromatography

Study of the retention behaviour of barbiturates by overpressured layer chromatography using silica gel bonded with tricaprylmethylammonium chloride
by K. Kovács-Hadady (Debrecen, Hungary) (Received August 9th, 1991) 301

Electrophoresis

Binary buffers for indirect absorption detection in capillary zone electrophoresis
by T. Wang and R. A. Hartwick (Binghamton, NY, USA) (Received September 11th, 1991) 307

Radial pH distribution during capillary electrophoresis with electroosomotic flow. Analysis with high ionic strength buffers
by A. Vinther (Gentofte and Lyngby, Denmark) and H. Søeberg (Lyngby, Denmark) (Received August 8th, 1991) . . 315

Capillary isoelectric focusing with electroosmotic zone displacement and on-column multichannel detection
by W. Thormann, J. Caslavska, S. Molteni and J. Chmelík (Berne, Switzerland) (Received September 12th, 1991) . . 321

Influence of poly(vinylpyrrolidone) on isotachophoretic separations of inorganic anions in aqueous electrolyte systems
by V. Madajová, E. Turcelová and D. Kaniansky (Bratislava, Czechoslovakia) (Received September 3rd, 1991) . . . 329

Separation of polycyclic aromatic hydrocarbons by micellar electrokinetic chromatography with cyclodextrins as modifiers
by Y. F. Yik, C. P. Ong, S. B. Khoo, H. K. Lee and S. F. Y. Li (Singapore) (Received August 8th, 1991) 333

Study of isotachophoretic separation behaviour of metal cations by means of particle-induced X-ray emission. II. Separation efficiency of lanthanoid model mixture
by J.-Y. Hu, T. Hirokawa, F. Nishiyama and Y. Kiso (Higashi-hiroshima, Japan) (Received August 13th, 1991) . . . 339

SHORT COMMUNICATIONS

Column Liquid Chromatography

Optical resolution of 2-chloro-3-phenylmethoxypropanoic acid after derivatization with (*S*)-2-octanol by high-performance liquid chromatography
by P. L. Anelli, C. Tomba and F. Uggeri (Milan, Italy) (Received September 10th, 1991) 346

Sensitive fluorescence detection of some nitrosamines by precolumn derivatization with dansyl chloride and high-performance liquid chromatography
by Z. Wang, H. Xu and C. Fu (Baoding, China) (Received September 24th, 1991) 349

Resolution of lipolysis mixtures from soybean oil by a solid-phase extraction procedure
W. E. Neff (Peoria, IL, USA), M. A. M. Zeitoun (Alexandria, Egypt) and D. Weisleder, Peoria, IL, USA) (Received July 30th, 1991) 353

Simultaneous ion chromatographic determination of chloride and calcium contained in Ringer's injection using a chelating agent as eluent
by W. Zhou, W. Liu and D. An (Jiangsu, China) (Received August 26th, 1991) 358

Solvent system for the counter-current purification of hydrophobic bombesin analogues
by L. Rusconi and M. Galantino (Milan, Italy) (Received June 6th, 1991) 362

Gas Chromatography

Study of dihydroxy-substituted saturated urushiol crown ether as a stationary phase in capillary gas chromatography
by Z.-R. Zeng, C.-Y. Wu, X.-H. Fang, Z.-F. Huang and Y.-T. Wang (Wuhan, China) (Received September 6th, 1991) 368

(Continued overleaf)

Direct gas chromatographic analysis of O,O-dimethyl- and O,O-diethyldithiophosphoric acids and identification of their by-products
by S. E. Massil, I. Ezra and L. Fuss (Beer-Sheva, Israel) (Received September 16th, 1991) 375

Monitoring 4-methoxy-2,3,6-trimethylbenzenesulphonyl deprotection of arginine-containing synthetic peptides using capillary zone electrophoresis
by F. Mulholland and G. R. Hague (Reading, UK) (Received October 11th, 1991) 380

Protection of terpenes against oxidative and acid decomposition on adsorbent cartridges
by A.-M. Strömvall and G. Petersson (Göteborg, Sweden) (Received September 30th, 1991) 385

Planar Chromatography

Determination of diacetoneketogulonic acid in water samples by high-performance thin-layer chromatography
by A. Eisenbeiss, S. Reuke and M. Türck (Darmstadt, Germany) (Received September 12th, 1991) 390

Author Index 395

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*).